

# The $\pi\pi$ mass spectrum in $Y(4260) \rightarrow \pi\pi J/\psi$ .

D.V. Bugg<sup>1</sup>

Queen Mary, University of London, London E1 4NS, UK

## Abstract

Three ways of fitting the  $\pi^+\pi^-$  mass spectrum in  $Y(4260)$  decays are studied. Data presented recently by Belle cannot be fitted by the  $\pi\pi$  S-wave intensity for elastic scattering. They can be fitted by adding a rather arbitrary destructive interference with the  $\sigma$  pole term. A better fit may be obtained with the decay sequence  $Y \rightarrow \pi H$ ,  $H \rightarrow \pi J/\psi$ , where  $H$  is a  $J^{PC} = 1^{--}$   $c\bar{c}q\bar{q}$  state peaking at 4.0 GeV with a width  $\sim 280$  MeV. A third possibility, involving a triangle diagram due to  $Y(4260) \rightarrow DD_1(2420)$ ,  $D_1 \rightarrow D^*(2007)\pi$ ,  $DD^* \rightarrow \pi J/\psi$  fails to fit the data. The first and second possibilities (or a combination) could be resolved by analysis of the  $Y(4260)$  Dalitz plot and  $\pi J/\psi$  mass projection, which are not presently publicly available.

PACS: 13.25.-k, 13.75.Lb, 14.40.Cs, 14.40.Ev.

There has been considerable interest in the  $Y(4260)$  discovered by BaBar [1]. It has been confirmed by Cleo [2], by further observations by BaBar [3,4] and Cleo [5] and most recently by Belle [6]. Its clear observation in decays to  $\pi\pi J/\psi$  has led to suggestions that it may be related to the  $c\bar{c}g$  hybrids predicted in this general mass range [7-10]. It is observed in the radiative return process, where it is produced through  $e^+e^- \rightarrow \gamma$ ; it therefore has  $J^{PC} = 1^{--}$ . Cleo observe  $Y(4260) \rightarrow \pi^0\pi^0 J/\psi$  [2] with a rate consistent with half that of  $\pi^+\pi^- J/\psi$ , albeit with just 8 clean events. This result strongly supports  $I = 0$  for  $Y(4260)$ .

The latest Belle data [6] reveal a curious  $\pi\pi$  mass spectrum. The objective here is to examine ways of fitting this spectrum. Three classes of decay scheme for  $Y(4260)$  are investigated.

The simplest hypothesis is  $Y \rightarrow (\pi\pi)_S J/\psi$ , where  $(\pi\pi)_S$  stands for the S-wave as observed in elastic scattering. The full curve of Fig. 1(a) compares data points with the spectrum predicted from the latest parametrisation [11]; a brief subroutine for this parametrisation is available from the author. In making this prediction, the line-shape of  $Y(4260)$  has been folded with the phase space for  $Y \rightarrow (\pi\pi)_S J/\psi$  using the Breit-Wigner parameters determined by Belle.

The predicted peak at  $\sim 900$  MeV is too wide, fails to reproduce the low point at 750 MeV and does not account for the low mass peak. A minor detail is that none of the fits reported here reproduce accurately the highest mass point at  $M(\pi\pi) \sim 1275$  MeV. This mass is outside the range accessible for Belle's central  $Y$  mass, 4295 MeV; it is possible that events in this bin arise from spreading of the high mass  $\pi\pi$  peak due to mass resolution. This bin is therefore omitted from all fits, and the curve is normalised to the 113 events in the remaining 9 bins. The resulting fit has  $\chi^2 = 69$  for 9 bins and 8 degrees of freedom.

In elastic scattering, the  $\pi\pi$  amplitude is

$$f_S(\pi\pi) = \frac{N(s)}{D(s)} = \frac{M\Gamma(s_{12})}{M^2 - s_{12} - iM\Gamma(s_{12})} \quad (1)$$

---

<sup>1</sup>email address: D.Bugg@rl.ac.uk




Figure 1: (a) The  $\pi\pi$  mass spectrum from Belle data for  $Y(4260) \rightarrow \pi^+\pi^- J/\psi$  (data points), compared with (i) production of  $J/\psi$  with the elastic  $\pi\pi$  S-wave (full curve), (ii) including destructive interference with the  $\sigma$  pole using eqn. (2) (dashed curve), (iii) with the triangle diagram of Fig. 3 (dotted); (b) the corresponding comparison with Cleo data.

where  $\Gamma(s)$  is close to linear in  $s$ , with a zero at the Adler point  $s = s_A = m_\pi^2/2$ . [The convention adopted here is that the  $J/\psi$  is taken as particle 3, hence  $s_{12}$  is the invariant mass of  $\pi\pi$ ]. The  $\pi\pi$  phase shift goes through  $90^\circ$  at  $M \sim 950$  MeV and the intensity of  $\pi\pi$  elastic scattering peaks there. However, the pole in the amplitude lies much lower, at  $\sim 470$  MeV, because of the strong  $s$ -dependence of the width. In some production processes, e.g.  $J/\psi \rightarrow \omega\pi\pi$  [12] and  $D \rightarrow 3\pi$  [13], the production amplitude is instead well fitted by the same Breit-Wigner denominator  $D(s)$  as eqn. (1), but with a constant numerator  $N(s)$ . The elastic amplitude seems to give a good fit to processes involving low momentum transfers, e.g. decays  $\psi(2S) \rightarrow \pi\pi J/\psi$  [14] and  $\phi \rightarrow \gamma\pi^0\pi^0$  [11]. The production processes of Refs. [12] and [13] involve larger momentum transfers, hence more distant singularities. The decay  $Y(4260) \rightarrow \pi\pi J/\psi$  is closer to the former case.

An adequate fit may be obtained using destructive interference between Eqn. (1) and the  $\sigma$  pole by altering the numerator  $N(s)$  to

$$f_S(\pi\pi) = \frac{(s - s_A) - G(M^2 - s_A)}{D(s)}, \quad (2)$$

with the same Breit-Wigner denominator  $D(s)$  as in eqn. (1). The resulting fit is shown by the dashed curve in Fig. 1, with  $G = 0.37$ ,  $\chi^2 = 17.8$  for 9 bins (7 fitted parameters). This fit seems somewhat artificial, since the numerator has no obvious explanation; however, it cannot presently be ruled out. Making  $G$  complex does not improve the fit significantly.

It seems unlikely that  $f_0(980)$  is responsible for the peak at 950 MeV in Fig. 1, because the  $f_0$  is too narrow. The latest BES data on  $J/\psi \rightarrow \phi\pi\pi$  and  $\phi KK$  [15] determine the full-width at half maximum of  $f_0(980) \rightarrow \pi\pi$  to be  $34 \pm 8$  MeV. Belle do not quote their  $\pi\pi$  mass resolution, but do quote the width of  $Y(4260)$  as  $133 \pm 26_{-6}^{+13}$  MeV. The mass resolution in both cases should be




Figure 2: (a) Fit to the Belle  $\pi\pi$  mass spectrum for  $L = 1$  decays via an intermediate  $H$  with  $M = 4080$  MeV,  $\Gamma = 280$  MeV; (b) the upper half of the Dalitz plot; (c) fit to Cleo data.

comparable. In any case, the  $\pi\pi$  mass distribution cannot be fitted by  $f_0(980)$  and the  $\sigma$  pole: the contribution from the  $\sigma$  pole is too broad if it interferes with only the very narrow  $f_0(980)$ . Cleo observe 2 events for the decay to  $K^+K^-J/\psi$  [2]. These could arise from the known amplitude for  $\sigma \rightarrow KK$  or from a small  $f_0(980)J/\psi$  contribution.

Fig. 1(b) compares the same fits with the 37 events of Cleo [2]. Although statistics are lower, the peak in data again looks narrower than the full curve; there is also some tentative indication of a rise at low  $\pi\pi$  mass. Cleo correct for background from sidebins around the  $J/\psi$ , but do not correct for possible background under the  $Y(4260)$  from other  $\psi$  states. Judging from the mass range 4.0 to 4.2 GeV, this background could be 15%.

BaBar [1] show a  $\pi\pi$  mass spectrum which does not show any rise at low  $\pi\pi$  mass. It has been obtained by summing the  $Y(4260)$  signal over the whole mass range and then making a background subtraction. The resulting signal/background ratio is  $\sim 1 : 1$ . In this respect, Belle's strategy of selecting events in the mass range 4.2 to 4.4 GeV, i.e. 1.5 times the full width, gives a better signal/background ratio for the purpose of viewing the  $\pi\pi$  mass spectrum. Some such weighting of data points near the peak of  $Y(4260)$  is desirable.

Let us turn to a second hypothesis: sequential decays of the form  $Y \rightarrow \pi H$ ,  $H \rightarrow \pi J/\psi$ . If  $Y$  has  $I = 0$ , as Cleo data suggest, the  $H$  has  $I = 1$  and could be a 4-quark state. Two cases need consideration. When the orbital angular momentum  $L$  in both steps of the sequence is zero,  $H$  has  $J^{PC} = 1^{+-}$ . If  $L = 1$ ,  $H$  has  $J^{PC} = 1^{--}$ ,  $0^{--}$  or  $2^{--}$ .

With  $L = 0$  in both steps, the predicted  $\pi\pi$  mass spectrum is close to the full curve of Fig. 1 and hence fails to fit the data. However, with  $L = 1$ , a good fit emerges naturally, as shown on Fig. 2(a). This result depends on simple kinematics. Consider the case  $Y_{123} \rightarrow \pi_1 H_{23}$ ,  $H_{23} \rightarrow \pi_2(J/\psi)_3$ , where numbers label the particles. If  $L = 1$  in the second step, there is a factor  $\cos\theta_2$  in the amplitude; here  $\theta_2$  is the angle of  $\pi_2$  in the 23 rest frame with respect to the recoil direction of particle 1. For fixed  $s_{23}$ ,  $s_{12}$  varies linearly with  $\cos\theta_2$  from one side of the Dalitz plot to the other, leading to a zero halfway between left- and right-hand edges of the plot. The mass of  $H$  determines the value of  $s_{12} = m_{12}^2$  at which the zero occurs, via the boundaries of the Dalitz plot. The width of  $H$  controls the sharpness of the dip in the mass spectrum near 750 MeV.

Fig. 2(b) shows the upper half of the Dalitz plot for this hypothesis. The zero due to the factor  $\cos \theta_2$  is clearly visible. Fig. 2(c) shows the fit to Cleo data. The fitted mass and width are almost linearly correlated because of the  $L = 1$  centrifugal barrier. The optimum fit has a mass of 4.08 GeV with  $\Gamma = 0.275$  GeV and  $\chi^2 = 15.1$  for 2 fitted parameters, seven degrees of freedom. Fits poorer in  $\chi^2$  by 1 are obtained with  $M = 4.04$  GeV,  $\Gamma = 0.225$  GeV or  $M = 4.16$  GeV,  $\Gamma = 0.330$  GeV. Because of the centrifugal barrier for production, the peak intensity is just below 4 GeV for all solutions. A  $Y$  mass at the lower end of the range is most plausible, in order to give a reasonable separation in masses of  $Y$  and  $H$ .

In detail, the amplitude for production of  $H_{23}$  alone is

$$f \propto f_Y(s_{123})f_H(s_{23})B_1(p_1)B_1(p_2)\cos\theta_2, \quad (3)$$

where  $f_Y$  is the Breit-Wigner amplitude of  $Y$  as a function of  $s_{123}$ ,  $f_H$  is the Breit-Wigner amplitude for  $H$ ,  $B_1(p_1)$  is the  $L = 1$  centrifugal barrier factor as a function of the momentum of  $\pi_1$  in the rest frame of  $Y$ , and  $B_1(p_2)$  is the centrifugal barrier factor as a function of the momentum of  $\pi_2$  in the rest frame of  $H_{23}$ . The centrifugal barrier is taken as  $B_1(p) = p/\sqrt{p^2 + p_0^2}$  where  $p_0 = 0.25$  GeV/c. The centrifugal barrier cuts off the intensity near the top of Fig. 2(b). The amplitude of eqn. (3) needs to be symmetrised between particles 1 and 2. The usual weighting according to phase space  $ds_{12}ds_{23}$  is also applied.

None of the experimental groups has presented data as a function of  $m^2(\pi J/\psi)$ . Even with the current low statistics, structure like that of Fig. 2(b) should be clearly visible. It gives rise to a strong peaking at two corners of the Dalitz plot and a peak in the  $\pi J/\psi$  spectrum near 4.0 GeV. The situation could however be confused by interferences between amplitudes of the two forms considered so far. Furthermore, interference between  $Y(4260)$  and underlying background from  $\psi(4160)$  and  $\psi(4415)$  is possible. Indeed, van Beveren and Rupp have argued against the existence of  $Y(4260)$  as a separate resonance [16]; they interpret the structure in this mass range as arising from thresholds and interferences in the  $s$ -dependence of  $\psi(4040)$  and  $\psi(4160)$  amplitudes.




Figure 3: The triangle diagram for  $Y \rightarrow DD_1(2420)$ ,  $D_1(2420) \rightarrow \pi D^*(2007)$ ,  $DD^*(2007) \rightarrow \pi J/\psi$ .

A third hypothesis will now be considered. The  $Y(4260)$  lies close to the threshold for  $D(1865)D_1(2420)$ , particularly for the resonance mass quoted by Belle, 4295 MeV. A possible

mechanism for production of the  $\pi\pi J/\psi$  system is shown in Fig. 3. In this so-called triangle diagram, the  $D_1$  decays (via a D-wave) to  $\pi D^*(2007)$ . This long-lived resonance can de-excite from the spectator  $D$  producing  $\pi J/\psi$ . A simulation of this process gives the dotted curve of Fig. 1(a), and fails to fit the Belle data. Even if interference with the  $(\pi\pi)_S J/\psi$  amplitude is included with a fitted complex coupling constant, nothing resembling the Belle mass spectrum can be reproduced. The essential feature of the process in Fig. 3 is that the  $D$  and  $D_1$  are produced almost at rest in the  $Y$  rest frame. The pion from  $D_1(2420)$  decay has a momentum close to 360 MeV/c in the  $D$  rest frame and hence in the  $Y$  rest frame. This makes a clear signature for this process.

In summary, the  $\pi\pi$  mass spectrum observed by Belle in  $Y(4260)$  decays is inconsistent with that expected for a final state  $(\pi\pi)_S J/\psi$  unless the numerator of the amplitude gives rise to destructive interference. The simplest, but speculative, hypothesis is the sequential decay to  $J/\psi$  via an intermediate  $J^{PC} = 1^{--}$  4-quark state. This possibility fits the data unaided; on the other hand, there is no obvious reason why the  $Y(4260)$  alone should prefer decays to  $\pi\pi J/\psi$ . It should be simple to verify or reject this hypothesis in its simplest form, although interferences with the first mechanism could complicate the situation.

## Acknowledgement

This work began during a stimulating visit to the Centro de Física das Interações Fundamentais of the Instituto Superior Técnico in Lisbon. I wish to thank Profs. E. van Beveren and G. Rupp for their hospitality and extensive discussions on this and related topics. I am grateful to Dr. F.-K. Guo for correcting a mistake in the original manuscript.

## References

- [1] B. Aubert et al. (BABAR collaboration), Phys. Rev. Lett. 95 (2005) 142001.
- [2] T.E. Coan et al. (CLEO collaboration), Phys. Rev. Lett. 96 (2006) 162003.
- [3] B. Aubert et al. Phys. Rev. D73 (2006) 012005.
- [4] B. Aubert et al. Phys. Rev. D73 (2006) 011101R.
- [5] Q. He et al. (CLEO collaboration), arXiv: hep-ex/ 0611021.
- [6] K. Abe et al. (BELLE collaboration), arXiv: hep-ex/ 0612006.
- [7] S.-L. Zhu, Phys. Lett. B625 (2005) 212.
- [8] F.E. Close and P.R. Page, Phys. Lett. B628 (2005) 215.
- [9] E. Kou and U. Pene, Phys. Lett. B631 (2005) 164.
- [10] X.Q. Luo and Y. Liu, Phys. Rev. D74 (2006) 034502.
- [11] D.V. Bugg, Eur. Phys. J C47 45.

- [12] M. Ablikim et al. (BES collaboration), Phys. Lett. B598 (2004) 149.
- [13] E.M. Aitala et al. (E791 collaboration), Phys. Rev. Lett. 86 (2001) 770.
- [14] M. Ablikim et al, arXiv: hep-ex/0610023.
- [15] M. Ablikim et al, Phys. Lett. B607(2004) 243.
- [16] E. van Beveren and G. Rupp, arXiv: hep-ph/0605317.